

UNIVERSITY OF BRISTOL

Summer 2025 Examination Period

FACULTY OF ENGINEERING

**Second Year Examination for the Degree of
Bachelor of Science and Master of Engineering**

**COMS20017
Algorithms & DATA**

**TIME ALLOWED:
2 Hours**

This paper contains *20* questions with a maximum of *100 marks*. Each question has exactly *one* correct answer. All answers will be used for assessment. You may use a non-programmable calculator.

The exam is closed-book (so no additional materials are allowed).

Blank pages are provided at the end for your workings. These working-outs will not be collected or marked. You must enter your answers on the provided answer sheet only. All questions in this examination will be computer marked. It is crucial that you follow the instructions and fill in the red coloured answer sheets carefully. Use a pencil (not pen) and a pencil eraser to correct any errors you make. Insert your candidate name, exam title, unit code and date on to the answer sheets in the relevant boxes. Fill in your student number.

Other Instructions:

TURN OVER ONLY WHEN TOLD TO START WRITING

Q1. What does the term 'Outlier' mean?

- A. A value that is left out of the analysis because of missing data
- B. When two data points are the same and one can be discarded
- C. An extreme value at either end of a distribution
- D. A type of variable that is not easy to quantify
- E. None of the above

[2 marks]

Q2. Given function $F(t) = \sum_{n=0}^8 12\cos(2\pi nt)$, what is the least number of sample points needed to satisfy the Nyquist criterion?

- A. 16
- B. 24
- C. 18
- D. 96
- E. 108

[4 marks]

Q3. Let the Fourier Series expansion of a function $f(x)$ to be:

$$f(x) = \sum_{n=0}^{\infty} a_n \cos\left(\frac{2\pi nx}{T}\right) + b_n \sin\left(\frac{2\pi nx}{T}\right).$$

Here are two different functions:

$$f_1(x) = x^2, \quad -\pi < x < \pi \quad \text{and} \quad f_2(x) = 1 + |x|, \quad -5 < x < 5$$

If functions $f_1(x)$ and $f_2(x)$ were expanded by their Fourier Series, then which is the correct statement below:

- A. All the a_n coefficients in both functions are zero.
- B. All the b_n coefficients in both functions are zero.
- C. The a_n coefficients in $f_1(x)$ and the b_n coefficients in $f_2(x)$ are zero.
- D. The b_n coefficients in $f_1(x)$ and the a_n coefficients in $f_2(x)$ are zero.
- E. For even n values only, the b_n coefficients in both functions are zero.

[5 marks]

Q4. Here are five statements about PCA (Principal Component Analysis):

1. PCA can be effective when multi-collinearity is suspected among features

(cont.)

2. After performing PCA in a 6D feature space and retaining the first three principal components, this means that the discarded components were not orthogonal to all other components.
3. The curse of dimensionality is when the number of features increases, so does the number of samples, resulting in a more elaborate model.
4. The purpose of eigenvectors in PCA is to determine the number of principal components to retain.
5. PCA is performed to reduce the number and increase the effectiveness of the features used.

Select the option below that is fully correct:

- A. 1, 2 and 3 are TRUE.
- B. 2, 3 and 4 are TRUE.
- C. Only 3 and 5 are TRUE.
- D. 3, 4 and 5 are TRUE.
- E. 1, 3 and 5 are TRUE.

[5 marks]

Q5. In the design of your autonomous driving system, you need to decide (as a binary classification problem) whether the path in front of the car shown by the front camera is 'road' or not. For this, you need to assess the performance of 3 different and independent features (f_a, f_b, f_c) by evaluating their t_{score} for 100 images taken by the front camera.

The mean and standard deviations of the resulting two classes for the 100 images and the size of each class (M and N) for each feature are:

$f_a : (\mu_1 = 3.5, \sigma_1 = 0.5), (\mu_2 = 1.5, \sigma_2 = 0.2), M = 75, N = 25$

$f_b : (\mu_1 = 12.2, \sigma_1 = 0.8), (\mu_2 = 5.7, \sigma_2 = 0.9), M = 67, N = 33$

$f_c : (\mu_1 = 1.5, \sigma_1 = 1.5), (\mu_2 = 4.4, \sigma_2 = 0.2), M = 78, N = 22$

Which of the following statements best explains the t_{score} values?

- A. f_a is the best feature, and f_b is the second best.
- B. f_b is the best feature, and f_c is the second best.
- C. f_a is the best feature, and f_c is the worst.
- D. f_b is the best feature, and f_a is the second best.
- E. f_c is the best feature, and f_b is the worst.

[8 marks]


Q6. The eigenvalues of the covariance of a dataset are: [26.0, 20.2, 14.3, 10.0, 7.8, 5.3, 2.0, 1.1, 0.77, 0.15]. How many eigenvalues represent 95.41% of the dataset?

(cont.)

- A. 5
- B. 6
- C. 7
- D. 8
- E. 9

[4 marks]

Q7. You are given a signal that needs to be sampled and quantised into binary values. You have already decided that the best sampling frequency is 4Hz and that you are going to use 3 bits to digitise the analog signal. What would be the correct binary representation of the signal plotted in the image below?



- A. 000011000010100011000111001
- B. 111010000001100011000001100
- C. 111011000010100011000011100
- D. 000011000010100111101111001
- E. 111011001010111000100111100

[6 marks]

Q8. Consider you are in Shanghai (at the origin of a coordinate system of distances to other cities). The coordinates of the city of Seoul relative to you is (80,115), while

(cont.)

the coordinates of the city of Manila relative to you is (10,-330). What is the Cosine Distance between Seoul and Manila?

- A. 1.803
- B. -0.803
- C. -0.197
- D. 0.197
- E. 1.197

[5 marks]

Q9. Here are a number of distances measured between two entities as stated:

- (i) $M = (-8,7)$, $N = (-2,-2)$ - Distance MN using ∞ -norm (L_∞) is 10
- (ii) $P = (5,6,12)$, $Q = (2,1,2)$ - Distance PQ using 3-norm (L_3) is 10.483
- (iii) $E = (-8,6,-5)$, $F = (-6,-4,9)$ - Distance EF using ∞ -norm (L_∞) is 4
- (iv) $W = \text{'Blasted'}$, $X = \text{'Plastered'}$ - Distance WX using Hamming Distance is 3
- (v) $Y = \text{'Divides'}$, $Z = \text{'Conquer'}$ - Distance YZ using Edit Distance is 6

Which of the results in the above statements are CORRECT:

- A. (ii), (iv) and (v).
- B. (i), (ii) and (iii).
- C. (iv) and (v)
- D. (i), (ii) and (iv).
- E. (ii) and (v).

[6 marks]

Q10. Your friend was recently on an 11-hour, long-haul flight. Each hour during the flight, he or she made a note of the "outside air temperature" reading as displayed on the aircraft information screen. The readings are shown as signal R below:

$$R = (0 \quad 25 \quad -44 \quad -70 \quad -70 \quad 50 \quad -40 \quad 0).$$

You decide to apply an averaging filter h to smooth this R signal, which then results in signal S (noting edge effects when performing convolution), i.e. $S = R * h$:

$$S = \frac{1}{6} (0 \quad 50 \quad -38 \quad -178 \quad -368 \quad -180 \quad -120 \quad 20 \quad -80 \quad 0).$$

Given $\frac{1}{6}$ as the normalisation factor, which of these filters below is the correct filter h ?

- A. $h = \frac{1}{6}(-1 \quad 4 \quad -1)$

(cont.)

- B. $h = \frac{1}{6}(2 \quad -2 \quad 2)$
- C. $h = \frac{1}{6}(-2 \quad 2 \quad -2)$
- D. $h = \frac{1}{6}(2 \quad 2 \quad 2)$
- E. $h = \frac{1}{6}(2 \quad 3 \quad 2)$

[5 marks]

Q11. Which of these statements about the natural logarithm and its use in data science is FALSE:

- A. The logarithm converts products into sums, i.e. $\ln x \cdot y = \ln x + \ln y$.
- B. The logarithm converts powers into products, i.e. $\ln x^a = a \ln x$
- C. The derivative of the natural logarithm is $\frac{\partial \ln x}{\partial x} = 2x^{-1}$
- D. When computing the Maximum Likelihood Estimate (MLE), the parameters with the highest log-likelihood are the same as the parameters with the highest “raw” likelihood.
- E. Using log-probabilities rather than “raw” probabilities helps us avoid numerical under/overflow.

[2 marks]

Q12. Which of the following statements about the MLE is FALSE?

- A. The MLE is an optimality criterion.
- B. The MLE is asymptotically unbiased.
- C. The MLE achieves the Cramer-Rao Lower Bound.
- D. The MLE is a consistent estimator.
- E. The MLE cannot always be obtained in closed form.

[3 marks]

Q13. Let x denote a Rayleigh distributed random variable with probability density function given by

$$p(x|\theta) = \frac{x}{\theta^2} \exp\left(-\frac{x^2}{2\theta^2}\right)$$

What is the MLE of θ ?

- A. $\hat{\theta}_{MLE} = \frac{1}{2} \sqrt{\frac{1}{N} \sum_{i=1}^N x_i^2}$
- B. $\hat{\theta}_{MLE} = \frac{1}{2N} \sqrt{\sum_{i=1}^N x_i^2}$
- C. $\hat{\theta}_{MLE} = \frac{1}{2N} \sum_{i=1}^N x_i^2$

(cont.)

D. $\hat{\theta}_{MLE} = \sqrt{\frac{1}{2N} \sum_{i=1}^N x_i^2}$

E. $\hat{\theta}_{MLE} = \frac{1}{N} \sum_{i=1}^N x_i^2$

[5 marks]

Q14. What is the estimator that minimizes the Bayes risk when the cost function used for its derivation is the absolute error between the estimate and its true value, i.e. $C[\theta, \hat{\theta}(x)] = C(\epsilon) = |\epsilon|$?

- A. The mode of the posterior distribution.
- B. The mode of the prior distribution.
- C. The median of the posterior distribution.
- D. The mean of the likelihood function.
- E. The median of the likelihood function.

[4 marks]

Q15. For a Maximum-a-Posteriori (MAP) estimator assume that the likelihood PDF is Gaussian with variance σ_n^2 defined as:

$$p(x|\theta) = \frac{1}{\sigma_n \sqrt{2\pi}} \exp \left[-\frac{(x - \theta)^2}{2\sigma_n^2} \right],$$

and that the prior pdf is a zero mean Gaussian with variance σ^2 defined as

$$p(\theta) = \frac{1}{\sigma \sqrt{2\pi}} \exp \left(-\frac{\theta^2}{2\sigma^2} \right).$$

What is the expression for the MAP estimate of θ ?

- A. $\hat{\theta} = \frac{\sigma_n^2}{\sigma^2} \cdot x$
- B. $\hat{\theta} = \frac{\sigma^2}{\sigma_n^2} \cdot x$
- C. $\hat{\theta} = \frac{\sigma^2}{\sigma_n^2 + \sigma^2} \cdot x$
- D. $\hat{\theta} = \frac{\sigma_n^2}{\sigma_n^2 + \sigma^2} \cdot x$
- E. A MAP estimate cannot be obtained in closed form.

[7 marks]

Q16. For the following data,

x	y
-2.1	0
-0.9	0
0.2	0
1.2	1
2.4	1

(cont.)

Fit a Gaussian distribution to each class, and compute the posterior probability that $x = 1.3$ is in class 1, given a prior of $P(y = 1) = 0.4$.

- A. 0.92
- B. 0.95
- C. 0.97
- D. 0.99
- E. 1.00

[8 marks]

Q17. In a two-class classification problem with a single feature, x , the pdfs are Gaussians with variances $\sigma^2 = 1/2$ for both classes and mean values 0 and 1, respectively:

$$P(x|\omega_1) = \frac{1}{\sqrt{\pi}} \exp(-x^2)$$

and

$$P(x|\omega_2) = \frac{1}{\sqrt{\pi}} \exp[-(x-1)^2]$$

If $P(\omega_1) = P(\omega_2)$ and the loss matrix is $L = \begin{pmatrix} 0 & 0.5 \\ 1 & 0 \end{pmatrix}$, what is the threshold value, x_0 , for minimum risk classification?

- A. $1/2$
- B. $1 + \ln 2/2$
- C. $(1 + \ln 2)/2$
- D. $(1 - \ln 2)/2$
- E. $1 - \ln 2/2$

[7 marks]

Q18. For the data displayed in the following table, compute the least-squares parameter fit for a model of the form $\hat{y} = w_1 + w_2x$

x	y
0.5	3.2
1.0	2.4
1.5	2.1
2.0	1.1

- A. $\hat{y} = 4.89 + 0.78x$
- B. $\hat{y} = 3.67 - 0.92x$
- C. $\hat{y} = 4.33 + 1.24x$
- D. $\hat{y} = 3.85 - 1.32x$

(cont.)

E. $\hat{y} = 3.85 - 1.36x$

[7 marks]

Q19. When is overfitting least likely to be a serious issue?

- A. When fitting a straight line (i.e. $y = w_0x + w_1$) with lots of training data.
- B. When little training data is available.
- C. When fitting a complex nonlinear function.
- D. when input data, $\mathbf{x}$, is a high-dimensional vector.
- E. When fitting a high-order polynomial.

[4 marks]

Q20. Which of the following statements is FALSE?

- A. Regularisation mitigates overfitting.
- B. Regularisation always improves prediction performance.
- C. Regularisation penalises very large values of the weights.
- D. The regularization parameter controls the strength of the penalty applied to the model's loss function.
- E. Regularization is particularly useful when fitting a complex function.

[3 marks]

The following pages are left blank for your rough workings. They will not be collected or marked. You must enter your answers on the provided answer sheet only.

